



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

It should be observed that the series converges only for values of x such that $x^3 - 3x > 2$. But $x^3 - 3x + 2 = (x + 1)^2(x - 2)$. Hence x must be greater than 2.

Also solved by A. G. CARIS, F. M. MORGAN, HORACE OLSON, ELIJAH SWIFT, S. W. REAVES, and T. M. BLAKESLEE.

402. Proposed by R. D. CARMICHAEL, Indiana University.

Obtain other series similar to that of Borda given in the preceding problem (No. 401).

SOLUTION BY S. W. REAVES, University of Oklahoma.

If in the well-known series

$$(1) \quad \log(n+1) = \log n + 2 \left[\frac{1}{2n+1} + \frac{1}{3} \left(\frac{1}{2n+1} \right)^3 + \frac{1}{5} \left(\frac{1}{2n+1} \right)^5 + \cdots \right]$$

we set

$$n = \frac{(x-2k)(x+k)^2}{4k^3},$$

and therefore

$$n+1 = \frac{(x+2k)(x-k)^2}{4k^3}, \quad \text{and} \quad \frac{1}{2n+1} = \frac{2k^3}{x^3-3k^2x},$$

we shall have a generalized Borda's series

$$(2) \quad \begin{aligned} \log(x+2k) &= 2 \log(x+k) - 2 \log(x-k) + \log(x-2k) \\ &+ 2 \left[\frac{2k^3}{x^3-3k^2x} + \frac{1}{3} \left(\frac{2k^3}{x^3-3k^2x} \right)^3 + \frac{1}{5} \left(\frac{2k^3}{x^3-3k^2x} \right)^5 + \cdots \right]. \end{aligned}$$

For $k=1$, (2) becomes Borda's series.

Again, if in (1) we set

$$n = \frac{(x-3k)(x+k)^3}{16k^3x},$$

we shall have

$$(3) \quad \begin{aligned} \log(x+3k) &= 3 \log(x+k) - 3 \log(x-k) + \log(x-3k) \\ &+ 2 \left[\frac{8k^3x}{x^4-6k^2x^2-3k^4} + \frac{1}{3} \left(\frac{8k^3x}{x^4-6k^2x^2-3k^4} \right)^3 + \cdots \right]. \end{aligned}$$

Also solved by F. M. MORGAN, T. M. BLAKESLEE, and A. M. HARDING.

GEOMETRY.

410. Proposed by A. H. HOLMES, Brunswick, Maine.

Given a focus and two tangents to an ellipse, prove that the locus of the foot of the normal corresponding to either tangent is a straight line.

SOLUTION BY A. M. HARDING, University of Arkansas.

Let SP and SP' be the fixed tangents, F the fixed focus, F' the other focus, and N the foot of the normal corresponding to SP . Draw FE and $F'E'$ perpendicular to SP , and FG and $F'G'$ perpendicular to SP' . Draw FF' cutting SP in L .